

Nonclassical paths in the recurrence spectrum of diamagnetic atoms

Brian E. Granger

ITAMP, Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Chris H. Greene

JILA and the Department of Physics, University of Colorado, Boulder, CO 80309-0440

Using time-independent scattering matrices, we study how the effects of nonclassical paths on the recurrence spectra of diamagnetic atoms can be extracted from purely quantal calculations. This study reveals an intimate relationship between two types of nonclassical paths: exotic ghost orbits and diffractive orbits. This relationship proves to be a previously unrecognized reason for the success of semiclassical theories, like closed-orbit theory, and permits a comprehensive reformulation of the semiclassical theory that elucidates its convergence properties.

INTRODUCTION

Recurrence spectroscopy [1] provides an excellent window into atomic systems whose classical counterparts are chaotic, such as atoms in strong magnetic fields. While the energy domain photoabsorption spectra of these systems are incredibly complex [2], the time domain physics is relatively simple, at least for short times. Recurrence spectroscopy gives access to this simple physics by revealing sharp (recurrence) peaks in the Fourier transform of the photoabsorption spectra. Semiclassical closed-orbit theory [3] provides an elegant interpretation of these short time features [4] in the spectra: each recurrence peak is associated with a classical trajectory in which the electron is launched from and returns to the nucleus radially. When the properties of these “closed orbits” (periods, stabilities, and so on) are used in the context of semiclassical approximations, the recurrence spectrum of hydrogen in a strong magnetic field can be predicted accurately [3, 5, 6].

There are, however, a number of limitations of closed-orbit theory. In its traditional formulation, it neglects quantum mechanical (Feynman) paths that have no classical counterparts. Examples of such paths are ghost orbits [7] (orbits that have not yet bifurcated into classical existence) and diffractive orbits [8] (orbits or paths that form when the electron scatters from features smaller than its wavelength). Clearly, fully quantum mechanical methods do not suffer from these shortcomings. A number of large scale quantum calculations have been performed on single channel atoms in strong magnetic fields [9]. While these calculations show excellent agreement with experiments they provide limited physical insight because they yield only the total photoabsorption or recurrence cross section, and do not necessarily help in its interpretation.

In this paper we outline a method for analyzing recurrence spectra that combines the interpretive strengths of closed-orbit theory with the ability of fully quantum mechanical methods to accurately describe nonclassical effects. This development relies on an exact quantum me-

chanical framework for treating atomic photoabsorption in external fields, which we derived in a previous paper [10]. When accurate quantum calculations are used in this framework, insight can be gained that is not available in either semiclassical methods or previous quantum calculations. To illustrate the power of this approach, we revisit the recurrence spectrum of diamagnetic hydrogen and study recurrences associated with diffractive and ghost orbits. Most significantly, an intimate relationship between these two types of orbits emerges. We conclude by discussing the nontrivial implications of this discovery for semiclassical theories of photoabsorption. Following others [5], we use the scaled variables (in atomic units) $w = 2\pi B^{-1/3}$ and $\epsilon = EB^{-2/3}$, so that in this paper the energy-like variable is the scaled field w and the time-like variable is the scaled action $\tilde{S}$.

These advances are enabled by our use of time-independent scattering matrices to describe the Rydberg electron’s motion exactly. When an external field is applied to an atom there are two such scattering matrices. First, the core-region S -matrix $\underline{S}^{\text{core}}$ of quantum defect theory [11] is used to characterize the scattering of the electron off the ionic core at small distances ($r < 10$ a.u.). For single channel atoms, such as the alkali-metal atoms, $\underline{S}^{\text{core}}$ is given in terms of the quantum defects μ_l : $S_{ll'}^{\text{core}} = \delta_{ll'} e^{2\pi i \mu_l}$.

Likewise, the long-range scattering matrix, $\underline{S}^{\text{LR}}$, characterizes the Rydberg electron’s motion in the combined Coulomb and diamagnetic potentials at distances far ($r > 10$ a.u.) from the ionic core. As closed-orbit theory first showed, it is both appropriate and fruitful to use semiclassical approximations for the electronic motion in this region. The semiclassical approximation to $\underline{S}^{\text{LR}}$ (which we do not utilize in any of the calculations in this paper),

$$S_{ll'}^{\text{LR}, \text{scl}}(w) = \sum_{j \in \text{co}} A_{ll'}^j(w) e^{i \tilde{S}_j w}, \quad (1)$$

is derived in [10]. Like closed-orbit theory, this involves a sum over the classical closed orbits having scaled actions $\tilde{S}_j$. The matrix $A^j(w)$ varies slowly with w and contains

other information about the closed orbits (Maslov index, stability, and so on). This form of the approximate, semiclassical $\underline{S}^{\text{LR}}$ suggests that the exact quantum mechanical $\underline{S}^{\text{LR}}$ can be written in a similar form [16], but with the sum being over all Feynman paths of the Rydberg electron that originate near the nucleus with angular momentum l' , and return to the nucleus with angular momentum l after scattering from the long range Coulomb and magnetic fields. While some of these quantum paths have classical analogues (closed orbits) also appearing in the semiclassical theory, others are purely quantal paths that are neglected in semiclassical theories.

The physics contained in the S -matrices can be probed with the photoabsorption cross section and its Fourier transform. The scattering matrices determine the accurate quantal photoabsorption cross section $\sigma(w)$ in a straightforward manner [10]:

$$\sigma(w) = 4\pi^2\alpha \text{Re}\omega_0 \vec{d} \left[\underline{1} - \underline{S}^{\text{core}} \underline{S}^{\text{LR}}(w) \right]^{-1} \times \left[\underline{1} + \underline{S}^{\text{core}} \underline{S}^{\text{LR}}(w) \right] \vec{d}^\dagger. \quad (2)$$

In this equation, ω_0 is the frequency of the light used to excite the transition and $\vec{d}$ is the atomic dipole vector. Equation 2 gives the energy smoothed cross section according to the following prescription: when the long-range S -matrix $\underline{S}^{\text{LR}}$ is calculated [12] at a complex value of the scaled field $w + i\frac{\Gamma}{2}$, Eq. 2 gives the total cross section “preconvolved” with a Lorentzian of width Γ [13].

The physical insight contained in the photoabsorption cross section can be extracted by expanding the matrix $\left[\underline{1} - \underline{S}^{\text{core}} \underline{S}^{\text{LR}} \right]^{-1}$ as a geometric series:

$$\sigma(w) \sim \text{Re} \vec{d} \left[\underline{1} + 2 \sum_{n=1}^{\infty} \left(\underline{S}^{\text{core}} \underline{S}^{\text{LR}} \right)^n \right] \vec{d}^\dagger. \quad (3)$$

Because the scattering matrices are simply quantum-mechanical amplitudes to scatter through a region of space a single time, the cross section can be viewed as an infinite sum over quantum mechanical paths where the Rydberg electron scatters one $\left(\underline{S}^{\text{core}} \underline{S}^{\text{LR}} \right)$ or more $\left(\underline{S}^{\text{core}} \underline{S}^{\text{LR}} \right)^n$ times from the core and long-range regions.

While the semiclassical $\underline{S}^{\text{LR},\text{scl}}$ (1) can be used in (3) to derive a semiclassical theory [10] of photoabsorption that resembles closed-orbit theory, we emphasize that it is not necessary to make a semiclassical approximation. In fact, by using an accurate, fully quantum mechanical $\underline{S}^{\text{LR}}$, detailed information about the recurrence spectrum can be extracted that is not available in either closed-orbit theory or other fully quantum mechanical methods. More specifically, we assert that the study of the Fourier transforms of individual terms in the expansion

of the cross section (3) reveals recurrences that appear only at a given order (n) of scattering. This ability to study only particular parts of the recurrence spectrum without making any semiclassical approximation is useful for studying recurrences associated with nonclassical paths. To illustrate how this analysis works, we now use it to study nonclassical paths in diamagnetic hydrogen without performing the detailed analysis required to repair the semiclassical approach [7].

There are two main types of nonclassical paths in diamagnetic atoms: ghost orbits and diffractive orbits. Because ghost orbits and diffractive orbits appear at different orders in the expansion of the photoabsorption cross section, their contributions can be isolated. This analysis is presented in Figs. 1, 2 and 4 where we present accurate quantum calculations of the scaled recurrence spectrum of diamagnetic hydrogen. These results were obtained by using an $\underline{S}^{\text{LR}}(w)$ calculated with a B -spline R -matrix technique and $\underline{S}^{\text{core}} = \underline{1}$ in the various terms of Eq. 3 [12]. The dipole vector $\vec{d}$ from the $2p$, $m = 0$ state to the even parity $m = 0$ final states shown here is given in [3]. In the scaled variables used here the energy-like variable, w , is related to an effective Planck’s constant $\hbar_{\text{eff}} = 2\pi/w$. With this in mind, we study the Fourier transform of the cross section $\sigma(w)$ at relatively low values of w (100 – 500) so that nonclassical effects are more pronounced.

The first order recurrence spectrum $\left| \text{FT} \left[2\text{Re} \vec{d} \underline{S}^{\text{LR}}(w) \vec{d}^\dagger \right] \right|$, pictured in Fig. 1, shows recurrence peaks from only primitive closed orbits and ghost orbits. To identify which recurrence peaks correspond to ghost orbits, we first perform a classical calculation of all the relevant closed-orbits at the scaled energy of interest. Then, when a first order recurrence peak is seen at a scaled action where no closed orbit exists, a ghost orbit has been found. More specifically, as Fig. 1 shows, we can readily identify the so called “exotic” ghost orbits that bifurcate into real closed orbits through a saddle-node bifurcation at (much) higher scaled energies. A semiclassical theory for these ghosts has been worked out by Main and Wunner [7] who predict an exponential falloff of the ghost amplitude at scaled energies below the classical bifurcation point. The first order recurrence spectrum in Fig. 1 clearly does not show such a decay of the ghost orbit recurrences. This apparent discrepancy, and its resolution, will be discussed below, after the second order recurrence spectrum has been presented.

The Fourier transform of the second order term in the expansion of the cross section, $\left| \text{FT} \left[2\text{Re} \vec{d} \underline{S}^{\text{LR}}(w) \underline{S}^{\text{LR}}(w) \vec{d}^\dagger \right] \right|$, is pictured in Fig. 2 and shows only recurrences due to double repetitions ($2R_1, 2V_1^1, \dots$) of primitive closed orbits and double combinations ($R_1 + V_1, \dots$) of different closed orbits. While the presence of recurrences due to repetitions




Figure 1: The Fourier transform, or recurrence strength, of the linear term in the expansion of the photoabsorption cross section $2\text{Re} \vec{d} \underline{S}^{\text{LR}}(w) \vec{d}^\dagger$ is plotted for hydrogen (even parity, $m = 0$) at seven scaled energies ($\epsilon = -0.9 \rightarrow -0.3$). Recurrence peaks corresponding to primitive closed orbits are labeled using the notation of [1]. Because an accurate, fully quantum mechanical $\underline{S}^{\text{LR}}(w)$ has been used, recurrence peaks associated with ghost orbits can also be seen. The recurrence peaks indicated by dashed lines are ghost orbits (X 's) [7] that bifurcate into real closed orbits through saddle-node bifurcations at higher scaled energies than shown here.

of the same closed orbit are predicted in semiclassical theories, the presence of combinations of *different* orbits in hydrogen is surprising from a semiclassical perspective. Typically it is argued (see [14], p.185) that because scattering of a low- l state by a pure Coulomb potential is strongly backscattered, the quantum wave returning to the nucleus along a given closed orbit can only scatter back out into the same closed orbit and not into any other closed orbits. However, we emphasize that this picture is strictly true only in the classical limit ($\hbar_{\text{eff}} = 0$). At any finite value of $\hbar_{\text{eff}}$, the $\Delta\theta\Delta l \gtrsim \hbar$ uncertainty relation limits the validity of this highly directional classical scattering argument. The accurate quantum calculations presented in Fig. 2 support this claim by showing second-order recurrences that appear at the scaled actions of combinations of primitive closed orbits.

The main point of the present letter is based on a simple observation about the first and second order recurrence spectra shown here. A quick comparison of Figs. 1 and 2 reveals an interesting trend about the recurrences of (exotic) ghost orbits and diffractive combination orbits. Namely, they have similar, if not identical, scaled actions and recurrence amplitudes. This correspondence holds for every case that we have studied, which suggests that there is a one to one relationship between exotic ghost orbits and diffractive combination orbits. This proposal is further elucidated by the topologies of the trajectories of the ghost and combination orbits, which are plotted in Fig. 3. *In all cases, orbits having similar scaled actions are also topologically similar.* We now turn to the most interesting aspect of this relationship, which




Figure 2: The recurrence strength, of the quadratic term in the expansion of the photoabsorption cross section $2\text{Re} \vec{d} \left(\underline{S}^{\text{core}} \underline{S}^{\text{LR}}(w) \right)^2 \vec{d}^\dagger$ is plotted for diamagnetic hydrogen (see Fig. 1). No semiclassical approximations have been made, which results the appearance of nonclassical recurrences that are related to the electron scattering twice from the long range region. More specifically, recurrences associated with combinations ($R_1 + V_1^1, \dots$) of two closed orbits are seen, which are classically forbidden because of the strong backscattering of classical low- l Coulomb scattering. Classically-allowed double repetitions of the primitive closed orbits ($2R_1, 2V_1^1, \dots$) are also seen.




Figure 3: This figure shows the topological similarity between the exotic ghost orbits (bottom) and their combination orbit counterparts (top). The orbits are plotted in cylindrical (ρ, z) coordinates at a scaled energy slightly above that where the ghost orbit has bifurcated into existence (as a true closed orbit) through a saddle-node bifurcation.

emerges when the total recurrence spectra (all orders of scattering) are plotted.

Surprisingly, the total recurrence spectra, $|\text{FT}[\sigma(w)]|$, shows neither exotic ghost orbits, nor combination orbits. More specifically, the ghost orbit recurrences (first order) cancel with the combination orbit recurrences (second order) in the total recurrence spectrum for hydrogen photoabsorption. Again, except very near the classical bifurcation points, this cancellation seems universal, and essentially complete.

This relationship between exotic ghost orbits and combination orbits in hydrogen can be expressed formally when the exact quantum $\underline{S}^{\text{LR}}$ is written in the form,

$$\underline{S}^{\text{LR}} = \sum_{j \in \text{co}} \underline{S}_j + \sum_{p \in \text{ghosts}} \underline{S}_p, \quad (4)$$




Figure 4: The recurrence strength of the total photoabsorption cross section $\sigma(w)$, Eq. (2), is plotted for diamagnetic hydrogen (see Figs. 1 and 2). While all nonclassical effects (ghost orbits, combination orbits) are included exactly in this purely quantal calculation, neither the ghost orbits (Fig. 1) nor the combination orbits (Fig. 2) survive in the total cross section. They cancel accurately.

where the first sum is over all paths associated with a classical closed orbit, and the second sum is over the ghost orbits that are not associated with a classically-allowed real closed orbit at the scaled energy of interest. Based on our accurate quantum calculations presented above, we postulate the following: *for every exotic ghost p there exist two or more primitive closed orbits $(i, j, k \dots)$ such that*

$$\underline{S}_p + P(\underline{S}_i \underline{S}_j \underline{S}_k \dots) \sim e^{-(\epsilon_b - \epsilon)}, \quad (5)$$

when the current scaled energy ϵ is less than the scaled energy ϵ_b where the ghost orbit bifurcates into a real closed orbit. The function $P()$ in (5) denotes the sum of all possible permutations of the primitive orbit S -matrices. The exponentially decaying recurrence amplitude that is predicted by Main and Wunner actually consists of two pieces: a diffractive orbit and a ghost orbit.

We close by outlining the implications of the cancellation (in hydrogen) between exotic ghost and combination orbits for semiclassical theories. Most importantly, in atoms other than hydrogen, the interference between these two types of paths is no longer completely destructive. This is because the diffractive combination orbits scatter twice from the ionic core, picking up a phase $\exp(4i\pi\mu_l)$, while the exotic ghost orbits scatter only once from the ionic core during their trajectory acquiring a phase of $\exp(2i\pi\mu_l)$. The resulting nondestructive interference leads to the well known core-scattered recurrences in the total spectra.

The cancellation condition, Eq. (5), can be incorporated into the photoabsorption cross section, Eq. (3), to derive a semiclassical theory of photoabsorption for atoms other than hydrogen. The resulting semiclassical

photoabsorption cross section is [12],

$$\sigma_{scl} \sim \text{Re} \, \vec{d}^\dagger \left[\underline{1} + 2 \underline{S}^{\text{core}} \underline{S}_{scl}^{\text{LR}} \sum_{n=0}^{\infty} \left(\underline{T} \underline{S}_{scl}^{\text{LR}} \right)^n \right] \vec{d} \quad (6)$$

which is valid except near the classical bifurcation points. This is exactly the result that other researchers have derived in extending closed orbit theory to treat nonhydrogenic atoms [8, 15], recast in terms of scattering matrices. Most importantly, this form of the cross section is a power series in the matrix $\underline{T} = \underline{1} - \underline{S}^{\text{core}}$, which vanishes for hydrogen. We emphasize that the assumptions implicit in closed-orbit theory, namely the subtle relationship between exotic ghost orbits and diffractive orbits, have remained obscured until now. This work elucidates these subtleties and describes how a meaningful semiclassical approximation emerges out of a careful consideration of the exact quantum mechanics.

The main advantage in writing the result of closed orbit theory in terms of S -matrices is that it is easily recognizable as a geometric series, which can be resummed analytically,

$$\sigma_{scl} \sim \text{Re} \, \vec{d}^\dagger \left(\underline{1} + 2 \underline{S}^{\text{core}} \underline{S}_{scl}^{\text{LR}} \left[\underline{1} - \underline{T} \underline{S}_{scl}^{\text{LR}} \right]^{-1} \right) \vec{d}, \quad (7)$$

to include all orders of core-scattering automatically. Furthermore, the convergence properties of the series, Eq. 6, are well known: Eq. (6) converges absolutely to Eq. (7) when:

$$\left| (\underline{S}^{\text{core}} - \underline{1}) \underline{S}_{scl}^{\text{LR}} \right| < 1. \quad (8)$$

While this is a formal requirement that may or may not be satisfied in a given circumstance, Eqs. (6), (7) and (8) provide a well established framework for further study of the convergence properties of closed orbit theory.

This work was funded in part by the U.S. Department of Energy, Office of Science. Discussions with John Delos and John Shaw were helpful in this work.

-
- [1] A. Holle *et al.*, Phys. Rev. Lett. **56**, 2594 (1986); **61**, 161 (1988).
 - [2] C. Iu *et al.*, Phys. Rev. Lett. **66**, 145 (1991).
 - [3] M. L. Du and J. B. Delos, Phys. Rev. A **38**, 1896 (1988); **38**, 1913 (1988).
 - [4] M. A. Al-Laithy, P. F. O'Mahony, and K. T. Taylor, J. Phys. B **19**, L773 (1986).
 - [5] J. Main *et al.*, Phys. Rev. A **49**, 847 (1994).
 - [6] K. Karremans, W. Vassen, and W. Hogervorst, Phys. Rev. A **60**, 4764 (1999).
 - [7] F. Haake, Act. Phys. Pol. A **85**, 693 (1994); J. Main and G. Wunner, Phys. Rev A **55**, 1743 (1997).
 - [8] P. A. Dando, T. S. Monteiro, D. Delande, and K. T. Taylor, Phys. Rev. A **54**, 127 (1996); J. S. Hersch, M. R. Haggerty, and E. J. Heller, Phys. Rev. Lett. **83**, 5342

- (1999); L. Wirtz, J. Z. Tang, and J. Burgdörfer, Phys. Rev. B **56**, 7589 (1997).
- [9] C. W. Clark and K. T. Taylor, J. Phys B **15**, 1175 (1982); P. F. O'Mahony, Phys. Rev. Lett. **63**, 2653 (1989); D. Delande, A. Bommier, and J. C. Gay, *ibid.* **66**, 141 (1991); S. Watanabe and H. Komine, *ibid.* **67**, 3227 (1991).
- [10] B. E. Granger and C. H. Greene, Phys. Rev. A **62**, 012511 (2000).
- [11] M. J. Seaton, Rep. Prog. Phys. **46**, 167 (1983).
- [12] B. E. Granger, Ph.D. thesis, University of Colorado at Boulder (2001).
- [13] F. Robicheaux, Phys. Rev. A **48**, 4162 (1993).
- [14] M. Courtney and D. Kleppner, Phys. Rev. A **53**, 178 (1996).
- [15] J. A. Shaw and F. Robicheaux, Phys. Rev. A **58**, 3561 (1998).
- [16] The exact $\underline{S}^{\text{LR}}$ can be cast in the same form as the semiclassical approximation, Eq. 1, using a path integral formulation of the scattering amplitudes.